Wheat Breeding for the Improved Bread-making Quality Using PCR Based Markers of Glutenins

Zuzana KOCOURKOVÁ 1 , Jana BRADOVÁ 2 , Zuzana KOHUTOVÁ 3 , Lucie SLÁMOVÁ 3 , Pavel VEJL 3 and Pavel HORČIČKA 1

¹Plant Breeding Station Stupice, SELGEN, a.s., Stupice, Czech Republic; ²Crop Research Institute, Prague-Ruzyně, Czech Republic; ³Czech University of Life Sciences in Prague, Prague, Czech Republic

Abstract: The relation between high-molecular-weight (HMW) glutenin subunits and bread-making quality could enable selection for improved bread-making quality in early stages of breeding process. The composition of HMW glutenin subunits was investigated in F_2 and F_7 progenies derived from the cross between winter wheat varieties Sulamit and Clever. The presence of Glu-A1 (AxNull and Ax1), Glu-B1 (Bx6+By8 and Bx 17+By18) and Glu-D1 alleles (Dx S+Dy10 and Dx 2+Dy12) was monitored using a PCR based assay. Segregation of alleles corresponded with the theoretically assumed 1:2:1 Mendelian ratio in F_2 generation, however, the values of χ^2 -test in F_7 generation indicated a strong affection of allelic frequencies by the breeding process. Significant variation was also observed in Glu-1 score frequency between F_2 and F_7 generation. These changes were probably caused by deliberate phenotypic selection for important agronomical traits. SDS and Zeleny sedimentation tests, mixographic parameter breakdown and HMW glutenin composition were analyzed in F_7 to reveal the effects of different combinations of HMW glutenin alleles on the bread-making quality characters. The results showed statistically significant differences in the contribution of HMW glutenin alleles. In general, the alleles Ax1, Bx17+By18 and Dx5+Dy 10 can be considered as markers of good baking quality. The data presented in this paper suggest that heterozygous constitution may also have a positive effect on bread-making quality.

Keywords: bread-making quality; breeding; glutenin subunits; MAS; winter wheat

Breeding for improved bread-making quality is an important breeding goal. This character is associated to a large extent with the amount and composition of endosperm seed proteins in the wheat kernel. Especially, the high-molecular-weight (HMW) glutenin subunits are considered to be significantly related to the technological quality of common wheat (PAYNE & CORFIELD 1979; AHMAD 2000; SCHWARZ *et al.* 2004).

The HMW glutenin subunits are encoded by genes at the *Glu-A1*, *Glu-B1* and *Glu-D1* loci on the long arms of homoeologous chromosomes 1A, 1B and 1D (Payne *et al.* 1981). Two tightly linked genes are present at each *Glu-1* locus, one encoding a

subunit of higher and the other of lower molecular weight, designated as the x and y type subunit, respectively (Harberd et al. 1986). At the Glu-1A locus, the x subunit is occasionally expressed and the y subunit is always absent. The x and y subunits coded by the Glu-1B and Glu-1D loci are almost always expressed, that means the Glu-B1 y-subunit can be absent. The HMW glutenin alleles Ax1 and Ax2* at Glu-1A locus, Bx17+By18 and Bx7+By8 or By9 at Glu-1B locus and Dx5+Dy10 at Glu-1D locus are connected with stronger dough and better baking properties, whereas AxNull, Bx6+By8 and Dx2+Dy12 are associated with poor baking quality (Payne et al. 1987).

Wheat breeders have been selecting wheat varieties with superior bread-making quality using SDS and Zeleny sedimentation tests, mixographic evaluation, baking test and other methods. However, these methods are not useful in very early hybrid generations, because they require a large amount of kernels and tests and they are destructive to tested samples. It is advantageous that the developed PCR based markers of HMW glutenin subunits may give quick characteristics of breadmaking quality already in the early hybrid generations (GALE 2005).

The aim of this study was to describe the HMW glutenin allelic composition in $\rm F_2$ and $\rm F_7$ generations of the progeny derived from the cross between winter wheat varieties Sulamit and Clever, using a PCR based assay, and to find out changes in allelic frequencies in these generations. It was also aimed to reveal the effects of different HMW glutenin composition on different bread-making quality characters.

MATERIAL AND METHODS

Plant material

Hybrid material obtained from the cross between winter wheat varieties Sulamit and Clever, differing in the composition of HMW glutenin subunits, was subjected to these analyses. The elite (E) parental variety Sulamit is characterized by the following composition of glutenins: AxNull/Bx17+By18/Dx5+Dy10. The variety Clever of "A" baking quality class had the following composition: Ax1/Bx6+By8/

Dx2+Dy12 (Bradová & Šašek 2005). Varieties Sulamit and Clever were crossed in 1999 at the breeding station in Stupice, Selgen a.s. The obtained plant progenies were gradually reproduced to the F₇ generation in 2006. Plants were selected according to their phenotypical traits mainly for resistance to rusts and also for high grain yield and winter hardiness since F, generation. Randomly selected F₂ individuals were obtained from the seeds stored in the gene bank of the breeding station in 2006. The PCR analyses of F₂ and F₇ individuals and the quality tests of F₇ individuals were carried out in 2006. The number of PCR investigated individuals in F2 and F2 generation was 120 and 88, respectively. Bread-making evaluation was carried out on 69 F₇ lines.

DNA extraction and PCR conditions

Genomic DNA was isolated from 100 mg of leaf tissues of single plants by an isolation kit GenElute Plant Genomic DNA Kit (Sigma).

PCR analyses were performed in a Biometra T-Gradient thermal cycler with heated lid. 25 μ l of PCR reaction mixture contained: 50–100 ng of template DNA, 1 × PCR buffer, 2.0 mmol/l MgCl₂, 0.4 μ mol/l of each primer, 0.2 mmol/l of dNTP, 1U *Taq* polymerase (Fermentas). Temperature conditions for the PCR reactions were 94°C for 5 min followed by 35 cycles of 94°C for 1 min, annealing temperature (see Table 1) for 45 s and 72°C for 1 min, followed by a final extension of 72°C for 5 min.

The horizontal electrophoresis from Bio-Rad with a 2% agarose gel in $1 \times TBE$ buffer was used

Table 1. The set of allele specific primers for HMW glutenin genes

Gene at Glu 1	References	Туре	Forward and reverse primers (5' \rightarrow 3')	Annealing temperature (°C)	
AxNull	Lafiandra et al. (1997)	dominant	ACGTTCCCCTACAGGTACTA	60	
			TATCACTGGCTAGCCGACAA		
Ax1 and $Ax2*$	Lafiandra <i>et al.</i> (1997)	dominant	CCATCGAAATGGCTAAGCGG	60	
			GTCCAGAAGTTGGGAAGTGC		
Dx5	D'Ovidio and Anderson (1994)	dominant	GCC TAG CAA CCT TCA CAA TC	(2)	
			GAA ACC TGC TGC GGA CAA G	63	
<i>Dy10</i> or <i>Dy 12</i>	Ѕмітн <i>et al.</i> (1994)	codominant	GTT GGC CGG TCG GCT GCC ATG	63	
			TGG AGA AGT TGG ATA GTA CC		
<i>Bx6</i> or <i>Bx7</i> or <i>Bx17</i>	MA et al. (2003)	codominant	CGCAACAGCCAGGACAATT	50	
			AGAGTTCTATCACTGCCTGGT	58	

for the confirmation of amplified DNA. DNA was visualized according to the standard methodology using ethidium bromide and documented by the Gel Doc XR documentation system (Bio-Rad). The presence of the specific part of a gene was verified using standard O'GeneRuler DNA Ladder Mix (Fermentas). Table 1 shows sets of allele specific primers used for the identification of *Glu-1* genes encoding individual HMW glutenin subunits.

Bread-making quality evaluation

SDS sedimentation test: 5 g of whole wheat flour was analyzed for sedimentation in SDS (sodium dodecyl sulphate). After specified shaking and rest time the sedimentation volume was measured. The test was performed using the procedure described by standard ČSN 461021 (1998).

Zeleny sedimentation test: 3.2 g of flour was analyzed for sedimentation in a lactic acid solution with isopropanol in the presence of bromophenol blue. After specified shaking and rest time the sedimentation volume was determined according to standard ISO 5529 (2000).


Mixographic parameter breakdown, which is associated with the weakening of the dough, was obtained from mixographic evaluation using a Reomixer instrument (ReoLogica Instruments AB). Mixing properties of dough according to standard

AACC 54-40A (1995) were obtained from 10 g of flour.

Each analysis was carried out at least twice in seed samples of F_7 generation (harvest in 2006).

RESULTS AND DISCUSSION

The five pairs of primers in AS-PCR (Allele Specific PCR) were used for the detection of alleles encoding the HMW glutenin subunits in 120 F₂ and 88 F₇ individuals derived from the cross between the cultivars Sulamit and Clever. Two primer sets designated by Lafiandra et al. (1997) identified the allelic composition at the Glu-1A locus. A 920 bp amplified fragment (Figure 1a) is characteristic of AxNull allele, the marker of poor bread-making quality. The absence of this fragment in the evaluated genotype showed the presence of one of the two alleles Ax1 or $Ax2^*$, which are considered to be the markers of good bread-making quality. A PCR fragment of 1.5 kb proved the presence of *Ax1* allele from the parent Clever (Figure 1b). The pair of primers according to MA et al. (2003) was used to verify alleles for HMW glutenin subunits at the Glu-B1 locus (Figure 1c). The presence of the high baking quality marker, allele Bx17 (PAYNE et al. 1987) from the variety Sulamit, was verified with the fragment of 669 bp after visualisation on the agarose gel. A strong association was described







Figure 1. Electrophoregrams of the alleles Ax1 (a), Ax-Null (b), Bx (c), Dx5 (d) and Dy (e) for randomly selected F_2 plants (1–10) and parental varieties Clever (C) and Sulamit (S)

between allele Bx6 and poor bread-making quality (SCHWARZ et al. 2004). This allelic composition, after detection by the AS-PCR method, was determined on agarose gel by the presence of two fragments of approximately 680 bp and 870 bp. Allele Bx7 was confirmed on the agarose gel by two fragments of 630 bp and 766 bp. The presence of a 450 bp band showed the occurrence of Dx5 HMW subunit allele that is associated with improved baking quality (D'Ovidio & Anderson 1994). No amplified fragment of 450 bp (Figure 1d) may be considered as a marker of the presence of other alleles at Glu-D1 locus (e. g. Dx2, Dx3 or Dx4) that are associated with poor bread-making quality. This marker turned out to be useless because of its dominant character. Dominant markers cannot discriminate between heterozygous and homozygous individuals. Early hybrid generations of wheat are still heterozygous. Hence, it was essential to use codominant markers or combination of more than one dominant marker for the differentiation

between alleles during segregating generations. Using the primer combination designed by SMITH et al. (1994), a difference between alleles Dy10 and Dy12 was confirmed. PCR reaction provided a fragment of 576 bp for *Dy10* allele and of 612 bp for Dy12 allele (Figure 1e). WITKOWSKI et al. (2005) used the sodium the method of dodecyl sulphate-polyacrylamide gel electrophoresis (SDS PAGE) to study the inheritance of HMW glutenin subunits coded by the Glu-1A locus. Because of the lack of specific band for the AxNull allele, this method was not successful in distinguishing heterozygous genotypes. The primer sets used in this study reliably distinguished the complete HMW glutenin allelic composition at all studied loci and determined heterozygous genotypes. Furthermore, this approach can be applied before harvest, because analyses are done from leaves. The SDS PAGE method is limited to be applied to grain, hence the analysis cannot be performed before harvest.






Figure 2. Comparison of the frequency distribution of HMW glutenin alleles at Glu-1 loci in F_2 and F_7 generation




Figure 3. Relationship between the *Glu-1* score (Payne *et al.* 1987) value for groups of HMW glutenin lines and frequencies of plants in F_2 and F_7 generation

According to the Mendelian inheritance, 25% of the F₂ plants were expected to be homozygous for each *Glu-1* allele and 50% were expected to be heterozygous. With the increased proportion of homozygous plants due to the self-pollination only 1.56% of the F₇ plants were expected to be heterozygous and 49.22% homozygous for each allele. Sample frequency distributions in F2 and F₇ generations were compared with these theoretical frequency distributions using χ^2 -test. The results of χ^2 -test confirmed the hypothesis of supposed theoretical distribution in F2 generation and disproved this hypothesis in F₇ generation. χ^2 -test values and significance probabilities for 2 degrees of freedom are shown in Table 2. The ratio 1:2:1 gained in F2 generation can be considered as verification of well performed cross. The values of χ^2 -test for F_7 generation indicate a strong affecting of Mendelian inheritance for the evaluated genes by the breeding process.

The observed allelic composition is summarized in Figure 2. It has been shown that allelic variation in HMW glutenins tended to the higher frequencies of alleles with a negative effect on bread-making quality in F₇ generation. It might be explained by targeted selection for resistance to the rust diseases, including leaf rust Puccinia recondita. The Lr 37 gene, conferring resistance to leaf rust, is present in a small translocation from Aegilops ventricosa on a short arm of 2A chromosome (BŁASCZYK et al. 2004). Конитоvá (unpublished) detected the Lr37 using CAPS marker in both F₂ and F₇ populations of progeny derived from Sulamit and Clever parents. Variety Clever is the donor of gene Lr37. The observed ratio of F₂ generation agreed with the 1:2:1 theoretically expected ratio ($\chi^2 = 1.11$, P = 0.57). However, 94.5% of F₇ genotypes were found to carry gene Lr37 in dominant homozygous constitution, which implies improvement in resistance to the leaf rust as a consequence of deliberate

Table 2. Verification of Mendelian segregation at a 1:2:1 ratio in F_2 and F_7 generations using χ^2 -test (df = 2)

Value –	Glu-1A	Glu-1A locus		Glu-1B locus		Glu-1D locus	
	F_2	F_7	F_2	\mathbf{F}_7	\mathbb{F}_2	F_7	
χ^2	0.38	118.72	0.82	62.25	0	92.68	
P	0.90-0.75	0.005-0	0.75-0.50	0.005-0	1.00-0.995	0.005-0	

df - degrees of freedom




Figure 4. Means and 95.0 percent confidence intervals for three bread-making quality characters and groups of lines differing in the composition of alleles encoding HMW glutenins at *Glu-1A* locus

phenotypic selection. A high level of this resistance seems to be associated with the presence of alleles responsible for deterioration of baking quality. A similar negative correlation between the leaf rust resistance and baking quality was reported by Drijepondt *et al.* (1990).

The frequency distribution of HMW glutenin alleles could be also influenced by the selection for winter hardiness. WITKOWSKI *et al.* (2008) have found the lines containing AxNull allele to be more frost tolerant than the lines containing AxI or $Ax2^*$ alleles, which suggests a linkage between winter hardiness and Glu-1A alleles responsible for deterioration of bread-making quality. Moreover, the parental variety Sulamit possessing AxNull allele is known to be more frost tolerant than the parental variety Clever with AxI allele.

Glu-1 score value for all allelic groups was calculated according to Payne $et\ al.$ (1987). The value for heterozygous loci was calculated as an average of both alleles present at the locus. Significant differences were observed in Glu-1 score frequency between F_2 and F_7 generation (Figure 3). It was found that F_2

Glu-1 score was in 13.3% of plants higher than and in 8.3% of plants the same as in the parental variety Sulamit of superior grain quality. 55.8% of plants had Glu-1 score higher than and 6.7% the same as the parental variety Clever (6). Only 15.8% of F_2 plants had a lower value than both parental varieties. By contrast, in F_7 generation, no plant with higher Glu-1 score value than Sulamit was found; 18.2% of plants reached a higher value than and 14.8% the same value as Clever in F_7 generation. 67.1% of evaluated individuals from the F_7 generation had assigned Glu-1 score below 6 and of it 47.7% of plants had the lowest obtained score (4).

The data on variation and average performance of 69 F7 lines in three grain quality characters, in comparison with parental varieties, are presented in Table 3. A comparison of the groups of lines differing in the composition of HMW weight subunits is provided in Table 4. The analysis of the results of SDS test, Zeleny test and the mixographic parameter breakdown showed the unequal contribution of alleles encoding HMW glutenin subunits to the examined parameters of bread-

Table 3. Results of SDS and Zeleny sedimentation tests (in ml) and mixographic evaluation (in mixograph units) of 69 lines of F_7 generation in comparison with the parental varieties Sulamit and Clever

	SDS test	Zeleny test	Mixograph breakdown
Average	74.97	42.67	1.55
SD	6.34	6.68	0.39
<i>CV</i> (%)	8.45	15.66	25.04
Minimum value	64	31	0.78
Maximum value	89	60	2.34
Sulamit	93	53	0.11
Clever	81	38	1.47

SD – standard deviation, CV – coefficient of variation




Figure 5. Means and 95.0% confidence intervals for three bread-making quality characters and groups of lines differing in the composition of alleles encoding HMW glutenins at *Glu-1B* locus

making quality. Luo *et al.* (2001) reported a high variation in the quality parameters associated with the possession of a certain allele. Three groups of genotypes were compared. The first group of genotypes differed in alleles encoding HMW glutenin subunits at the Glu-1A locus. The second group had various alleles at the Glu-1B locus. No heterozygous genotype was found in F_7 generation for this group. The third group differed in HMW alleles at the Glu-1D locus.

Differences in the genotypic group variant in Glu-1A alleles were statistically significant (P < 0.05) for Zeleny test and mixographic parameter breakdown. Heterozygous allelic composition and Ax1 allele at

this locus showed a positive effect on Zeleny test. A positive effect on the mixographic parameter breakdown, which is in negative correlation with bread-making quality, was found for heterozygous allelic composition (not significant) and Ax1 allele (P < 0.05). No statistically significant difference was however found in the SDS sedimentation volume. These results correspond with the results of Payne $et\ al.\ (1987)$ or Nieto-Taladriz $et\ al.\ (1994)$. The data presented in this study suggest that heterozygous constitution may also have a positive effect on bread-making quality (Table 4, Figure 4). Significant differences (P < 0.05) between Null/17+18/2+12 and Null/6+8/2+12 allelic com-

Table 4. Means of bread-making quality parameters for groups of HMW glutenin lines in F_7 generation in comparison with parental varieties

Composition of HMW glutenin subunits	No. of lines	Payne score value	SDS test (ml)	Zeleny test (ml)	Mixograph breakdown (MU)
H 6+8 2+12	8	5.0	73.0a	46.1a	1.4a
N 6+8 2+12	32	4.0	73.7a	39.4b	1.9b
1 6+8 2+12	3	6.0	74.0a	46.7ab	1.5a
N 6+8 2+12	32	4.0	73.7a	39.4a	1.9a
N 17+18 2+12	10	6.0	83.0b	42.8a	1.3b
N 6+8 H	6	5.5	72.5a	45.5ab	1.2a
N 6+8 2+12	32	4.0	73.7a	39.4a	1.9b
N 6+8 5+10	10	7.0	73.9a	47.4b	1.1a
Sulamit (N 17+18 5+10)		9.0	93.0	53.0	0.1
Clever (1 6+8 2+12)		6.0	81.0	38.0	1.5

Different letters in a column denote a statistically significant difference of LSD test (P < 0.05); N – AxNull allele; H – heterozygous composition at the locus; MU – mixograph unit




Figure 6. Means and 95.0% confidence intervals for three bread-making quality characters and groups of lines differing in the composition of alleles encoding HMW glutenins at *Glu-1D* locus

position were found in SDS sedimentation volume and mixographic parameter breakdown. A positive influence of the lines with 17+18 subunits on both quality tests was confirmed. This result confirms previous observations of Peltonen *et al.* (1993). There was no significant difference in the means of Zeleny test between the genotypes differing in Bx6+By8 and Bx17+By18 alleles (Table 4, Figure 5). Dx5+Dy10 and heterozygous constitution at the Glu-1D locus had a positive effect (P < 0.05) on Zeleny test and mixographic parameter breakdown. A negative effect of Dx2+Dy12 was published previously (SHEWRY et al. 1992; AHMAD 2000; CAMPOS et al. 2004; TABIKI et al. 2006 and others) and our results confirm the results of these studies. There was no significant difference in the means of SDS test between the genotypes carrying Dx5+Dy10 and Dx2+Dy12 alleles. Heterozygous constitution at this locus showed a statistically significant (P < 0.05) positive influence on the breakdown parameter (Table 4, Figure 6).

Payne *et al.* (1987) developed a predictive scoring system for individual HMW glutenin subunits based on the SDS sedimentation volume. However, only the allelic pair Bx17+By18 showed a difference in the SDS volume (Table 4). It could probably be ascribed to differences in the genetic background or to a reduced variability of lines in F_7 generation for this trait as a consequence of phenotypic selection for resistance to the rusts. As shown in Table 3, the coefficient of variation in the SDS sedimentation test was the lowest (8.5%), when compared with Zeleny test (15.7%) and mixographic parameter breakdown (25.0%).

In general, the obtained results demonstrate the usefulness of application of PCR based markers of glutenins in wheat breeding programs. It can be expected that the determination of HMW glutenin subunits composition can significantly contribute to the improvement of bread-making quality already in early hybrid generations. However, it should be taken into consideration that a deliberate selection for e.g. Ax1 at Glu-A1 may result in lower winter hardiness (WITKOWSKI et al. 2008). With early strong selection for winter hardiness, or rust resistance in our case, the alleles with a negative impact on the end use quality can be fixed in advanced breeding materials.

Acknowledgements. Supported by Czech Science Foundation, Project No. 521/05/H013 and by the Czech University of Life Sciences in Prague, Project No. IG 20/2006.

References

AHMAD M. (2000): Molecular marker-assisted selection of HMW glutenin alleles related to wheat bread quality by PCR-generated DNA markers. Theoretical and Applied Genetics, **101**: 892–896.

BŁASCZYK L., GOYEAU H., HUANG X., RÖDER M., STĘPIEŃ Ł., СНЕŁKOWSKI J. (2004): Identifying leaf rust resistance gene *Lr37* on the microsatellite map of wheat. Cellular & Molecular Biology Letters, **9**: 869–878.

Bradová J., Šašek A. (2005): Diversity of gliadins and HMW glutenin subunits in Czech registered wheat varieties. Czech Journal of Genetics and Plant Breeding, **41**: 160–163.

CAMPOS H., ZUNIGA J., RATHGEB P., JOBET C. (2004): Selection of Chilean wheat genotypes carriers of the HMW glutenin allele *Glu-D1 x5* through polymerase chain reaction. Agricultura Técnica, **64**: 223–228.

D'OVIDIO R., ANDERSON O.D. (1994): PCR analysis to distinguish between alleles of member of a multigene family correlated with bread-making quality. Theoretical and Applied Genetics, **88**: 759–763.

- DRIJEPONDT S.C, PRETORIUS Z.A., VAN LILL D., RIJ-KENBERG F.H.J. (1990): Effect of *Lr34* resistance on leaf rust development, grain yield and baking quality in wheat. Plant Breeding, **105**: 62–68.
- GALE K.R. (2005): Diagnostic DNA markers for quality traits in wheat. Journal of Cereal Science, **41**: 181–192.
- Harberd N.P., Bartels D., Thompson R.D. (1986): DNA restriction fragment variation in the gene family encoding high-molecular-weight (HMW) glutenin subunits of wheat. Biochemical Genetics, **24**: 579–596.
- LAFIANDRA D., TUCCI G.F., PAVONI A., TURCHETTA T., MARGIOTTA B. (1997): PCR analysis of x- and y-type genes present at the complex *Glu-A1* locus in durum and bread wheat. Theoretical and Applied Genetics, **94**: 235–240.
- Luo C., Griffin W.B., Branlard G., McNeil D.L. (2001): Comparison of low and high molecular weight wheat glutenin allele effects on flour quality. Theoretical and Applied Genetics, **102**: 1088–1098.
- MA W., ZHANG W., GALE K.R. (2003): Multiplex-PCR typing of high molecular weight glutenin alleles in wheat. Euphytica, **134**: 51–60.
- NIETO-TALADRIZ M.T., PERRETANT M.R., ROUSSET M. (1994): Effect of gliadins and HMW and LMW subunits of glutenin in the $\rm F_6$ recombinant inbred lines from a bread wheat cross. Theoretical and Applied Genetics, $\bf 88$: 81–88.
- Payne P.I., Corfield K.G. (1979): Subunits composition of wheat glutenin proteins, isolated by gel filtration in dissociating medium. Planta, **145**: 83–88.
- Payne P.I., Holt L.M., Law C.M. (1981): Structural and genetical studies of the HMW subunits of the wheat glutenin. 1. Allelic variation in subunits amongst varieties of the wheat (*Triticum aestivum* L.). Theoretical and Applied Genetics, **60**: 229–236.
- Payne P.I., Nigtingale M.A., Krattiger A.F., Holt L.M. (1987): The relationship between HMW glutenin

- subunit composition and the bread-making quality of British-grown wheat varieties. Journal of the Science of Food and Agriculture, **40**: 51–65.
- Peltonen J., Salopelto J., Rita H. (1993): The optimal combination of HMW glutenin subunits coded at gene loci *Glu-A1* and *Glu-B1* for bread-making quality in Scandinavian wheats. Hereditas, **118**: 71–78.
- SHEWRY P.R., HALFORD N.G., TATHAM A.S. (1992): High-molecular weight subunits of wheat glutenin subunits of wheat glutenin. Journal of Cereal Science, 5: 105–120.
- SMITH R.L., SCHWEDER M.E., BARNETT R.D. (1994): Identification of glutenin alleles in wheat and triticale using PCR-generated DNA markers. Crop Science, **34**: 1373–1378.
- Schwarz G., Felsenstein F.G., Wenzel G. (2004): Development and validation of a PCR-based marker assay for negative selection of the HMW glutenin allele *Glu-B1-1d* (*Bx6*) in wheat. Theoretical and Applied Genetics, **109**: 1064–1069.
- Tabiki T., Ikeguschi S., Ikeda T.M. (2006): Effects of High-molecular-weight and Low-molecular-weight glutenin subunit alleles on common wheat flour quality. Breeding Science, **56**: 131–136.
- WITKOWSKI E., WAGA J., BIELAWSKA A., WITKOWSKA K., LUBER H. (2005): Inheritance of glutenin proteins coded by the locus on chromosome *1A* in F₂ hybrid genotypes of winter wheat. Biuletyn Instytutu Hodowli i Aklimatyzacji Roślin, **235**: 57–64.
- WITKOWSKI E., WAGA J., WITKOWSKA K., RAPACZ M., GUT M., BIELAWSKA A., LUBER H., LUKASZEWSKI A. J. (2008): Association between frost tolerance and the alleles of high molecular weight glutenin subunits present in Polish winter wheats. Euphytica, **159**: 377–384.

Received for publication May 14, 2008 Accepted after corrections September 17, 2008

Corresponding author:

Ing. Zuzana Kocourková, SELGEN, a.s., Šlechtitelská stanice Stupice, Stupice 24, 250 84 Sibřina, Česká republika

tel.: + 420 281 012 458, fax: + 420 281 012 466, e-mail: kocourkova@selgen.cz